

[Free Full Text from Publisher](#)[Look Up Full Text](#)

Save to EndNote online

[Add to Marked List](#)

5 of 100

PRODUCTION AND STABILITY OF MYCOFLOCCULANTS FROM LENTINUS SQUARROSULUS RWF5 AND SIMPLICILLIUM OBCLAVATUM RWF6 FOR REDUCTION OF WATER TURBIDITY

By: [Jebun, N](#) (Jebun, Nessa)^[1]; [Alam, MZ](#) (Alam, Md. Zahangir)^[1]; [Al-Mamun, A](#) (Al-Mamun, Abdullah)^[1]; [Raus, RA](#) (Raus, Raha Ahmad)^[1]

[View ResearcherID and ORCID](#)**IJUM ENGINEERING JOURNAL****Volume:** 19 **Issue:** 1 **Pages:** 48-58**DOI:** 10.31436/ijumej.v19i1.843**Published:** 2018**Document Type:** Article

Abstract

The production and stability of two novel myco-flocculants produced by river water fungus (RWF) were investigated. Screening tests were conducted to find suitable nutrients, pH, nutrient concentration, inoculum dose, and stability for two mycoflocculants *L. squarrosulus* (RWF5) and *S. obclavatum* (RWF6). The strains showed good flocculating activity in reducing turbidity of kaolin suspension while malt extract was used as nutrient source. Supernatants of RWF5 and RWF6 were able to reduce turbidity from 900 +/- 10 NTU to 46 NTU (95%) and 195 NTU (78%), respectively. In order to enhance the production, optimization of cultivation conditions were studied using a one-factor-at-a-time (OFAT) method. *L. squarrosulus* (RWF5) reduced 96% of turbidity at optimum conditions, comprising of 0.1% (w/v) malt extract, 3% (v/v) inoculum dose, and initial pH 7.0 for 6 days. The results of the compatible mixed culture showed good flocculation activity at 88% compared to a single culture of *S. obclavatum* at 78%. On the other hand, *L. squarrosulus* showed better turbidity reduction in the single culture rather than the mixed culture. The stability of *L. squarrosulus* and *S. obclavatum* supernatants showed excellent turbidity reduction over a wide pH range of 4-8 with the maximal flocculation rate of 96% and 90%, respectively, at pH 7.0. They also exhibited high turbidity removal ability in a temperature range of 4 C-degrees - 55 C-degrees for 24h with a maximum turbidity removal rate of 96% (RW5) and 87% (RW6) at 25 C-degrees. Time stability of the *L. squarrosulus* supernatant showed good turbidity removal potential at above 90% at room temperature (28 +/- 2 C-degrees) and 85% at low temperature (4 C-degrees) for 12 days. The high flocculating rate of the myco-flocculants and their good stability under wide range of temperature indicated their potentiality as biodegradable flocculants for water and wastewater treatment industry.

Keywords

Author Keywords: [flocculation](#); [jar test](#); [Lentinus squarrosulu](#); [mixed culture](#); [Simplicillium obclavatu](#); [stability](#); [turbidity](#)

KeyWords Plus: [WASTE-WATER](#); [MOLECULAR-IDENTIFICATION](#); [FLOCCULATION PROPERTIES](#); [BACILLUS-LICHENIFORMIS](#); [BIOFLOCCULANT](#); [SLUDGE](#); [STRAINS](#); [BIOCONVERSION](#); [POTENTIALITY](#); [PERFORMANCE](#)

Author Information

Reprint Address: Alam, MZ (reprint author)

Citation Network

In Web of Science Core Collection

0

Times Cited

[Create Citation Alert](#)

27

Cited References

[View Related Records](#)

Use in Web of Science

Web of Science Usage Count

0

Last 180 Days

0

Since 2013

[Learn more](#)

This record is from:

Web of Science Core Collection

- Emerging Sources Citation Index

Suggest a correction

If you would like to improve the quality of the data in this record, please [suggest a correction](#).

+ Int Islamic Univ Malaysia, Fac Engr, Dept Biotechnol Engr, BERC, Jalan Gombak, Kuala Lumpur 53100, Malaysia.

Addresses:

+ [1] Int Islamic Univ Malaysia, Fac Engr, Dept Biotechnol Engr, BERC, Jalan Gombak, Kuala Lumpur 53100, Malaysia

E-mail Addresses: zahangir@iium.edu.my

Funding

Funding Agency	Grant Number
Ministry of Higher Education (MOHE)	FRGS-14-109-0350
Research Management Centre (RMC), International Islamic University Malaysia	

[View funding text](#)

Publisher

KULLIYAH ENGINEERING, INT ISLAMIC UNIV MALAYSIA, JALAN GOMBAK, 53100, MALAYSIA

Categories / Classification

Research Areas: Engineering

Web of Science Categories: Engineering, Multidisciplinary

[See more data fields](#)

◀ 5 of 100 ▶

Cited References: 27

Showing 27 of 27 [View All in Cited References page](#)

(from Web of Science Core Collection)

- Evaluation of fungal potentiality for bioconversion of domestic wastewater sludge** **Times Cited: 19**
 By: Alam, MZ; 'I-Razi, AF; Molla, AH
 JOURNAL OF ENVIRONMENTAL SCIENCES Volume: 16 Issue: 1 Pages: 132-137 Article Number: 1001-0742(2004)16:1<132:EOFPFB>2.0.TX;2-H Published: 2004
- Enhanced settleability and dewaterability of fungal treated domestic wastewater sludge by liquid state bioconversion process** **Times Cited: 55**
 By: Alam, MZ; Fakhru'I-Razi, A
 WATER RESEARCH Volume: 37 Issue: 5 Pages: 1118-1124 Article Number: PII S0043-1354(02)00452-9 Published: MAR 2003
- Production of a bioflocculant from Aspergillus niger using palm oil mill effluent as carbon source** **Times Cited: 35**
 By: Aljuboori, Ahmad H. Rajab; Uemura, Yoshimitsu; Osman, Noridah Binti; et al.
 BIORESOURCE TECHNOLOGY Volume: 171 Pages: 66-70 Published: NOV 2014
- Production and characterization of a bioflocculant produced by Aspergillus flavus** **Times Cited: 53**
 By: Aljuboori, Ahmad H. Rajab; Idris, Azni; Abdullah, Norhafizah; et al.
 BIORESOURCE TECHNOLOGY Volume: 127 Pages: 489-493 Published: JAN 2013
- Flocculation behavior and mechanism of bioflocculant produced by Aspergillus flavus** **Times Cited: 20**
 By: Aljuboori, Ahmad H. Rajab; Idris, Azni; Al-joubory, Hamid Hussain Rijab; et al.
 JOURNAL OF ENVIRONMENTAL MANAGEMENT Volume: 150 Pages: 466-471 Published: MAR 1 2015